

## RESEARCH ARTICLE

# Effects of pre-treatments on bulk stable isotope ratios in fish samples: A cautionary note for studies comparisons

Maëlle Connan<sup>1</sup>  | Grant Hall<sup>2</sup>  | Malcolm Smale<sup>1,3</sup> 

<sup>1</sup>Institute for Coastal and Marine Research, Marine Apex Predator Research Unit, Department of Zoology, Nelson Mandela University, PO Box 77000, Port Elizabeth 6031, South Africa

<sup>2</sup>UP Stable Isotope Laboratory, Mammal Research Institute, University of Pretoria, Private Bag X20, Hatfield 0028, South Africa

<sup>3</sup>Port Elizabeth Museum, PO Box 13147 Humewood, Port Elizabeth 6013, South Africa

**Correspondence**

M. Connan, Institute for Coastal and Marine Research, Marine Apex Predator Research Unit, Department of Zoology, Nelson Mandela University, PO Box 77000, Port Elizabeth 6031, South Africa.

Email: maelle.connan@gmail.com

**Funding information**

African Coelacanth Ecosystem Programme of the National Research Foundation, Grant/Award Number: 81879

**Rationale:** Stable isotope analysis (SIA) has revolutionised ecological studies over the past thirty years. One of the major fields where SIA is applied in the marine environment is related to the definition of ecosystem structure and function. With marine top predators such as sharks, SIA is a method of choice because tissue samples can be collected without the sacrifice of the animal. In elasmobranch research, the influence of compounds such as urea, trimethylamine oxide and lipids must be considered when using stable isotopes as ecological markers. Currently, a range of pre-treatments are used to chemically remove these molecules prior to SIA.

**Methods:** This study investigated the impact of eleven commonly used pre-treatments on carbon and nitrogen contents and C:N atomic ratio, as well as carbon and nitrogen SI ratios in elasmobranch tissues and its prey, measured by isotope ratio mass spectrometry. Three tissues were tested: blood and muscle of the ragged-tooth shark *Carcharias taurus*, and muscle of one teleost species, the Cape knifejaw *Oplegnathus conwayi*.

**Results:** Compared with untreated samples, no trend or generalisation could be highlighted with the influence of pre-treatments being species-, tissue- and chemical-element-dependent. For the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values, differences among pre-treatments were as high as 3‰, therefore potentially leading to erroneous ecological interpretation.

**Conclusions:** The chemical properties of compounds (e.g. urea, lipids) combined with the polarity of solutions (e.g. water, solvents) explained a large part of these observations. This study highlights that pre-treatments need to be considered especially when comparing carbon and nitrogen stable isotope ratios between studies. The results of this study provide a call to all stable isotope researchers to make a concerted effort to standardise pre-treatment methods. This is crucial as global reviews are becoming increasingly more informative.

## 1 | INTRODUCTION

The use of biomarkers to study the trophic ecology of predators has enhanced tremendously the understanding of such interactions in the last 30 years. A favoured method is the measurement of naturally occurring stable isotopes in an animal's tissues. In the marine environment, measurements of carbon ( $^{13}\text{C}/^{12}\text{C}$ ;  $\delta^{13}\text{C}$  values) and nitrogen ( $^{15}\text{N}/^{14}\text{N}$ ;  $\delta^{15}\text{N}$  values) stable isotope ratios

are the most common indicators of carbon flow pathways and consumer's trophic position, respectively.<sup>1</sup> The stable isotope ratios of these elements in a consumer's tissues result from an equilibrium between diet and physiology.<sup>2</sup>

Similarly to other fields, fish scientists are increasingly using stable isotopes as spatial and trophic indicators and studies have increased tremendously over the last 10 years, bringing new insights into fish trophic ecology (e.g.<sup>3</sup>), migration patterns (e.g.<sup>4</sup>), and physiology

(e.g.<sup>5</sup>). Studies of the foraging ecology of elasmobranchs are no exception with a recent increase in published papers using the stable isotope approach (e.g.<sup>6-9</sup>). Due to the life history of elasmobranchs, their wide-ranging migration patterns and concern about their population decline,<sup>10</sup> the stable isotope approach for their study is particularly useful, as samples can be collected non-lethally (e.g. biopsies), and by sampling various tissues of different metabolic rates from the same individual, trophic information can be obtained at various time scales (e.g. muscle vs blood<sup>11,12</sup>). However, the elevated urea ((NH<sub>2</sub>)<sub>2</sub>CO) and trimethylamine *N*-oxide (C<sub>3</sub>H<sub>9</sub>NO; TMAO) concentrations in elasmobranch tissues used to combat osmotic stress<sup>13-15</sup> complicate the interpretation of stable isotope ratios when used as ecological indicators. Typically, urea is an end product of nitrogen metabolism which is enriched in <sup>14</sup>N.<sup>16</sup> Tissues with high urea content will thus exhibit lower  $\delta^{15}\text{N}$  values and C:N ratios than those for the same tissue types with low urea contents. This factor is particularly important for lipid-rich tissues because lipids are known to artificially increase C:N ratios and lower  $\delta^{13}\text{C}$  values.<sup>17-20</sup> Tissues rich in both lipids and urea may therefore exhibit completely acceptable C:N ratios when in fact their  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values are affected by urea and lipid contents, respectively.<sup>21</sup>

Numerous different protocols are currently being used to chemically remove urea and lipids from elasmobranch tissues. To date, little is known about how these various protocols compare in their efficiency of lipid and urea removal. This is likely to affect carbon and nitrogen contents and  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values, as well as C:N ratios. Critically important information for comparisons between studies (e.g.<sup>22</sup>) is thus missing. To investigate this, we tested the influence of 11 pre-treatments commonly used in published studies on carbon and nitrogen percentages and C:N atomic ratios, as well as carbon and nitrogen stable isotope ratios, from muscle tissue for two fish species: one elasmobranch species, the ragged-tooth shark (*Carcharias taurus* Rafinesque 1810), and one teleost species, the Cape knifejaw (*Oplegnathus conwayi* Richardson 1840). In addition, we tested whether the influence of pre-treatments was tissue-dependent by comparing treated muscle and blood samples from individual ragged-tooth sharks. Due to their different biochemical compositions, we hypothesised that the influence of pre-treatments on stable isotope ratios and C:N ratios would be species- and tissue-specific.

## 2 | EXPERIMENTAL

### 2.1 | Sample collection

Muscle samples of Cape knifejaw ( $n = 5$ ), and whole blood and muscle samples of ragged-tooth sharks ( $n = 8$ ) were collected between March 2012 and October 2014 along the south-east coast of South Africa. Cape knifejaw samples were collected from dead fish (stranded individuals killed by an anoxic phytoplankton bloom), as part of a larger study looking at the ecological role of sharks in the area. Ragged-tooth shark samples were mostly collected from living animals that were captured in an ultrasonic tagging study to investigate their movement pattern. When sharks were in tonic immobility, muscle tissue was collected with a corer to remove a tissue sample ventral to the dorsal

fin, and blood was collected from the caudal vein using a disposable syringe and needle. Sampled ragged-tooth shark were released within 10 min of capture once all samples had been collected. Two of the ragged-tooth sharks were sampled freshly dead (one stranded and one caught as bycatch by local commercial fisheries). Samples were stored on ice in the field and then frozen to  $-20^{\circ}\text{C}$  on return to the laboratory. They were later oven-dried at  $50^{\circ}\text{C}$  for 48 h. Samples were then finely ground prior to further analysis.

### 2.2 | Sampling

The homogenised samples were sub-divided into seven sub-samples before being subjected to a range of pre-treatment methods to remove urea and lipids from the various tissue types (Figure 1).


Set 1 of the samples was left untreated and are referred to as *Bulk*. The other six sample sets were treated using a range of lipid and urea extraction methods commonly used in fish and elasmobranch studies (e.g.<sup>23-25</sup>).

Set 2 was treated using repeated washes with distilled water (referred to as H<sub>2</sub>O) to remove urea from the various tissues, based on the methodology used by Li et al.<sup>25</sup> and Kim and Koch.<sup>23</sup> Aliquots of tissue were placed in 5-mL graduated centrifuge tubes and 4 mL of distilled water added and vortexed (IKA MS 3 basic; IKA®-Werke, Staufen, Germany) for 1 min. They were placed on a mixer (Coulter Electronics Ltd, Harpenden, UK) and rotated for 24 h, centrifuged at 5000 rpm for 5 min and the water removed with a syringe. This was repeated a further two times and the samples were then dried at  $50^{\circ}\text{C}$ .

Set 3 was treated with petroleum ether, the solvent used by Kim and Koch.<sup>23</sup> Aliquots of the samples were placed in 1.5-mL micro-centrifuge tubes with 1.2 mL of petroleum ether (referred to as *PE*; Minema P1630; Spellbound Laboratory Solutions, Port Elizabeth, South Africa) and vortexed for 1 min, placed on the Coulter mixer for 1 h, centrifuged at 5000 rpm for 5 min and the petroleum ether decanted. This was repeated a second time and the samples were dried at  $50^{\circ}\text{C}$ .

Set 4 was treated with cyclohexane, the solvent used by Kiszka et al.<sup>24</sup> Aliquots of the samples were placed in 1.5-mL micro-centrifuge tubes with 1.2 mL of cyclohexane (referred to as *Cx*; SAAR1763000LC; UniLab, Merck (Pty) Ltd, Modderfontein, South Africa) and vortexed for 1 min, placed on the Coulter mixer for 1 h, centrifuged at 5000 rpm for 5 min and the cyclohexane decanted. This was repeated a second time and the samples were then dried at  $50^{\circ}\text{C}$ .

Sets 5 and 6 were both treated with a 2:1 chloroform/methanol mixture adapted from Hussey et al.<sup>26-28</sup> Lipid extraction on sample set 5 was performed using a single rinse of the 2:1 chloroform/methanol solution (referred to as *CM1*; chloroform: SAAR1595040LC uniVar; methanol: SAAR4146080LC uniVar; Merck (Pty) Ltd, Modderfontein, South Africa), while set 6 received two rinses (referred to as *CM2*). Aliquots of sample were placed in 1.5-mL micro-centrifuge tubes with 1.2 mL of the 2:1 chloroform/methanol solution and vortexed for 1 min, placed on the Coulter mixer for 1 h, centrifuged at 5000 rpm for 5 min and the chloroform/methanol decanted. This was repeated a second time for sample set 6 and the samples were then dried at  $50^{\circ}\text{C}$ .



**FIGURE 1** Schematic representation of the various pre-treatments tested. SI: stable isotope

Set 7 was pre-treated in a similar manner to *CM1*, but using a 2:1 chloroform/ethanol solution (referred to as *CEth*; ethanol rectified 96% SAAR2233510LP uniVar, Merck (Pty) Ltd) with the samples receiving a single rinse. This is currently the lipid extraction method used by the Mammal Research Institute Stable Isotope Laboratory at the University of Pretoria (South Africa).<sup>29</sup>

Five additional pre-treatments were then prepared to test the effect of additional water rinses on the solvent washed tissues as in Li et al.<sup>25</sup> Solvent-washed samples *PE*, *Cx*, *CM1*, *CM2* and *CEth* were subjected to a series of distilled water washes over a period of 3 days as per the method used for the *H<sub>2</sub>O* samples. These water-treated samples are referred to as *PE H<sub>2</sub>O*, *Cx H<sub>2</sub>O*, *CM1 H<sub>2</sub>O*, *CM2 H<sub>2</sub>O* and *CEth H<sub>2</sub>O*, respectively (Figure 1).

### 2.3 | Stable isotope analysis

All sub-samples of ragged-tooth shark whole blood and muscle, and Cape knifejaw muscle (*Bulk*, *H<sub>2</sub>O*, *PE*, *Cx*, *CM1*, *CM2*, *CEth*, *PE H<sub>2</sub>O*, *Cx H<sub>2</sub>O*, *CM1 H<sub>2</sub>O*, *CM2 H<sub>2</sub>O*, *CEth H<sub>2</sub>O*) were then analysed for carbon and nitrogen stable isotopic composition. Aliquots of 0.55 to 0.60 mg of each sub-sample were weighed into tin capsules which

had been pre-cleaned in toluene (SAAR6081040LC uniVar, Merck (Pty) Ltd).

Several samples were run in duplicate to ensure reproducibility of the results. Isotopic analysis was carried out by continuous-flow isotope ratio mass spectrometry via an elemental analyser (Flash EA 1112 Series) coupled to a Delta V Plus stable light isotope mass spectrometer via a ConFlo IV system (all equipment supplied by Thermo Fisher, Bremen, Germany), housed at the Stable Isotope Laboratory, Mammal Research Institute, University of Pretoria, Pretoria, South Africa.

Two in-house laboratory running standards, Merck Gel ( $\delta^{13}\text{C} = -20.57\text{‰}$ ,  $\%C = 43.83$ ,  $\delta^{15}\text{N} = 6.80\text{‰}$ ,  $\%N = 14.64$ ) and DL-valine ( $\delta^{13}\text{C} = -10.57\text{‰}$ ,  $\%C = 41.28$ ,  $\delta^{15}\text{N} = -6.15\text{‰}$ ,  $\%N = 15.29$ ), as well as a blank sample, were run after every 11 unknown samples. The analytical accuracy of these laboratory running standards was validated against international standards (National Institute of Standards & Technology; NIST, Gaithersburg, MD, USA): NIST 1557b (bovine liver), NIST 2976 (mussel tissue) and NIST 1547 (peach leaves). All samples were measured and calibrated using the above-mentioned laboratory running standards. The carbon and nitrogen percentages and atomic C:N ratios for each sample were calculated using a mass balance equation utilising the sample and Merck Gel weights, and the known

C (41.28%) and N (15.29%) percentages for the Merck Gel. Each run comprised 96 unknown samples, 20 laboratory standards and 10 blanks. All results are referenced to the internationally used standards of V-PDB (Vienna Pee-dee Belemnite) for carbon isotope ratios, and to atmospheric air for nitrogen isotope ratios. Results are expressed in delta ( $\delta$ ) notation using the standard equation:

$$\delta X (\text{‰}) = (R_{\text{sample}}/R_{\text{standard}}) - 1 \quad (1)$$

where  $X = {}^{13}\text{C}$  or  ${}^{15}\text{N}$  and  $R$  represents  ${}^{13}\text{C}/{}^{12}\text{C}$  or  ${}^{15}\text{N}/{}^{14}\text{N}$ .

The analytical precision, based on the results for the DL-valine standard, across multiple runs was  $<0.1\text{‰}$  for both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values.

## 2.4 | Statistical analysis

The effect of pre-treatments on carbon and nitrogen percentages, C:N atomic ratios and carbon and nitrogen stable isotope ratios was approached by calculating the difference between treated tissues and *Bulk* for all three sample types. Either a paired Student's t-test or Wilcoxon signed rank tests were then used to assess whether the differences observed were significant. Paired Student's t-tests were used when normality (Shapiro test) and homoscedasticity (Levene's test) assumptions were verified. Differences among pre-treatments were then tested separately on each tissue using analyses of variance (ANOVAs) for repeated sampling when residuals followed a normal

distribution. When this assumption was not verified, data were rank transformed and repeated ANOVAs were then conducted. Both were followed by Tukey post-hoc tests using a Bonferroni correction factor.

The level of significance was set at 0.05 or less in the case of Bonferroni correction for multiple comparisons. All statistical analyses were conducted using PAST v3.0<sup>30</sup> and R<sup>31</sup> software.

## 2.5 | Ethics


This work was undertaken under an annual research permit issued by the South African Department of Agriculture, Forestry and Fisheries and Department of Environmental Affairs: Oceans and Coasts (DEA: O&C). Animal handling was according to the Standard Operating Procedure: Shark Tagging, accepted by Bayworld and DEA: O&C Ethics Committees.

## 3 | RESULTS

### 3.1 | Elemental composition

The water rinsing of *Bulk* tissues had mixed effects on carbon and nitrogen percentages as well as C:N ratios depending on tissue and species (Figure 2).

Rinsing of Cape knifejaw muscle did not change the carbon ( $48.3 \pm 3.5\%$  vs  $48.3 \pm 0.4\%$ ) or nitrogen ( $13.5 \pm 0.1\%$  vs  $13.2 \pm 0.8\%$ )



**FIGURE 2** Changes in carbon and nitrogen percentages (Perc. C and Perc. N, respectively) and C:N ratios after the eleven pre-treatments relative to the untreated *Bulk* samples (Cape knifejaw muscle, ragged-tooth shark muscle and blood). The data for water-rinsed tissues are in blue. (Refer to Figure 1 for abbreviations) [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

percentages but significantly increased the C:N ratios ( $4.2 \pm 0.1$  vs  $4.3 \pm 0.1$ ; Table 1a). For ragged-tooth shark muscle, water rinsing significantly lowered nitrogen percentage ( $14.9 \pm 0.9\%$  vs  $13.7 \pm 1.2\%$ ) and increased C:N ratios ( $2.9 \pm 0.1$  vs  $4.1 \pm 0.1$ ) (Table 1b). The water treatment on untreated samples (*Bulk*) significantly affected the three parameters for ragged-tooth shark blood (increased % C:  $36.9 \pm 2.8\%$  vs  $50.6 \pm 0.6\%$  and C:N atom:  $2.5 \pm 0.3$  vs  $4.3 \pm 0.1$ , or decreased % N:  $17.1 \pm 0.7\%$  vs  $13.7 \pm 0.8\%$ ) (Table 1c).

Compared with *Bulk*, no trend or generalisation could be highlighted as the influence of pre-treatments was species-, tissue- and chemical-element-dependent (Table 2). The tissue most affected by pre-treatments was ragged-tooth shark blood with 26 out of 30 outcomes of comparisons being significant (Table 2c).

Most subsequent water rinses of tissues pre-treated with solvents did not affect the carbon and nitrogen percentages in Cape knifejaw muscle samples except for the *CM1*, *CM2* and *CEth* pre-treatments (6 out of 10 outcomes of the comparisons were significant; Table 2a, Figure 3). In contrast, water rinses on solvent pre-treated samples significantly affected most carbon and nitrogen contents in both ragged-tooth shark tissues except for muscle nitrogen content with the *CEth* pre-treatment (Tables 2b and 2c, Figure 3). The C:N ratios were diversely significantly affected depending on pre-treatments and tissues (Table 2, Figure 3).

The final values of carbon and nitrogen percentages as well as C:N ratios were significantly affected by pre-treatments (all three ANOVAs on ranks  $p < 0.001$ ; Table 3). Most outcomes of the pair-wise comparisons were significant for the three tissues (Tables S1–S3, supporting information).

### 3.2 | Stable isotopic composition

The water rinse of *Bulk* tissues did not significantly affect the  $\delta^{13}\text{C}$  values for muscle tissue of either Cape knifejaw ( $-15.4 \pm 0.2\%$  vs  $-15.2 \pm 0.3\%$ ) or ragged-tooth shark ( $-14.3 \pm 0.9\%$  vs  $-14.2 \pm 0.7\%$ ), as opposed to the  $\delta^{15}\text{N}$  values which significantly increased in muscle tissue samples of both Cape knifejaw ( $12.8 \pm 0.4\%$  vs  $13.7 \pm 0.3\%$ ) and ragged-tooth shark ( $14.8 \pm 0.7\%$  vs  $15.5 \pm 0.7\%$ ; Tables 1a and 1b, Figure 4). Both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values decreased significantly in ragged-tooth shark blood after water rinsing compared with untreated *Bulk* samples ( $\delta^{13}\text{C}$ :  $-13.2 \pm 0.7\%$  vs  $-14.1 \pm 0.6\%$ ;  $\delta^{15}\text{N}$ :  $15.0 \pm 0.7\%$  vs  $13.7 \pm 0.8\%$ ; Table 1c).

The water rinsing after solvent wash significantly affected  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in Cape knifejaw muscle and ragged-tooth shark blood (except for the *PE* pre-treatment of Cape knifejaw muscle; Table 2). The direction of the effect (increased or decreased) was, however, species-, tissue- and chemical-element-dependent (Figure 3). For ragged-tooth shark muscle, the water rinsing also significantly increased  $\delta^{15}\text{N}$  values compared with tissues pre-treated with solvents, but  $\delta^{13}\text{C}$  values were only affected by water rinsing after the pre-treatments *PE*, *CM2* and *CEth* (Table 2, Figures 3 and 4).


Pre-treatments had significant effects on the final  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in all three tissues (Table 3, Figure 4). For muscle samples from both species, the difference in  $\delta^{13}\text{C}$  values among all pre-treatments added up to 0.7‰, while the differences in  $\delta^{15}\text{N}$  values added up to 1.6‰ and 1.8‰ for ragged-tooth shark and Cape knifejaw, respectively (Figure 4). For ragged-tooth shark blood, the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  ranges were 1.8‰ and 1.7‰, respectively (Figure 4).

**TABLE 1** Effect of the water rinsing on untreated samples (*Bulk*) or subsequent to solvent washing on carbon and nitrogen percentages, C:N ratios, and  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in the three fish tissues. (Refer to Figure 1 for abbreviations). Significant results are in bold, W values are in italics

	<b>H<sub>2</sub>O</b>		<b>PE</b>		<b>Cx</b>		<b>CM1</b>		<b>CM2</b>		<b>CEth</b>	
	t or W	p	t or W	p	t or W	p	t or W	p	t or W	p	t or W	p
<i>a) Cape knifejaw muscle (n = 5)</i>												
Percent C	-0.01	0.995	-1.80	0.146	15	0.063	-3.98	<b>0.016</b>	15	0.063	-18.28	<b>&lt;0.001</b>
Percent N	0.64	0.556	-0.99	0.379	15	0.063	-8.97	<b>&lt;0.001</b>	15	0.063	-17.21	<b>&lt;0.001</b>
C:N atomic	-6.78	<b>0.002</b>	-5.24	<b>0.006</b>	12	0.313	27.5	<b>&lt;0.001</b>	4.15	<b>0.014</b>	-12.51	<b>&lt;0.001</b>
$\delta^{13}\text{C}$	-1.93	0.126	6.93	<b>0.002</b>	3.33	<b>0.029</b>	-6.19	<b>0.003</b>	-6.08	<b>0.004</b>	12.25	<b>&lt;0.001</b>
$\delta^{15}\text{N}$	-18.62	<b>&lt;0.001</b>	-18.21	<b>&lt;0.001</b>	-3.97	<b>0.017</b>	-29.07	<b>&lt;0.001</b>	-5.89	<b>0.004</b>	-7.22	<b>0.002</b>
<i>b) Ragged-tooth shark muscle (n = 8)</i>												
Percent C	-2.33	0.052	-6.49	<b>&lt;0.001</b>	36	<b>0.012</b>	36	<b>0.012</b>	36	<b>0.012</b>	-8.85	<b>&lt;0.001</b>
Percent N	33	<b>0.036</b>	5.12	<b>0.002</b>	4.7	<b>0.002</b>	36	<b>0.012</b>	-5.05	<b>0.002</b>	28	0.161
C:N atomic	36	<b>0.012</b>	28	<b>0.018</b>	-8.11	<b>&lt;0.001</b>	36	<b>0.012</b>	27.5	0.184	36	<b>0.012</b>
$\delta^{13}\text{C}$	-0.85	0.421	4.09	<b>0.006</b>	2.36	0.051	-0.89	0.403	-4.85	<b>0.002</b>	35	<b>0.017</b>
$\delta^{15}\text{N}$	-6.35	<b>&lt;0.001</b>	-6.02	<b>&lt;0.001</b>	-4.68	<b>0.002</b>	-17.22	<b>&lt;0.001</b>	-8.58	<b>&lt;0.001</b>	-2.80	<b>0.026</b>
<i>c) Ragged-tooth shark blood (n = 8)</i>												
Percent C	36	<b>0.012</b>	36	<b>0.012</b>	36	<b>0.012</b>	35	<b>0.017</b>	-6.17	<b>&lt;0.001</b>	36	<b>0.012</b>
Percent N	36	<b>0.012</b>	36	<b>0.012</b>	5.96	<b>&lt;0.001</b>	-3.99	<b>0.005</b>	-6.66	<b>&lt;0.001</b>	-3.17	<b>0.016</b>
C:N atomic	36	<b>0.012</b>	36	<b>0.012</b>	36	<b>0.012</b>	0.96	0.366	0.42	0.685	-8.81	<b>&lt;0.001</b>
$\delta^{13}\text{C}$	16.4	<b>&lt;0.001</b>	14.73	<b>&lt;0.001</b>	9.85	<b>&lt;0.001</b>	-6.99	<b>&lt;0.001</b>	-11.69	<b>&lt;0.001</b>	3.08	<b>0.018</b>
$\delta^{15}\text{N}$	7.1	<b>&lt;0.001</b>	8.17	<b>&lt;0.001</b>	8.07	<b>&lt;0.001</b>	-12.52	<b>&lt;0.001</b>	-3.08	<b>0.018</b>	3.25	<b>0.014</b>

**TABLE 2** Changes in carbon and nitrogen percentages, C:N ratios, and  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values from the 10 pre-treatments relative to untreated (Bulk) muscle (Cape knifejaw, ragged-tooth shark) and blood (ragged-tooth shark) samples. (Refer to Figure 1 for abbreviations). Significant results are in bold, W values are in italics

	PE		PE H <sub>2</sub> O		Cx		Cx H <sub>2</sub> O		CM1		CM1 H <sub>2</sub> O		CM2		CM2 H <sub>2</sub> O		CEth		CEth H <sub>2</sub> O			
	t	W	p	t	W	p	t	W	p	t	W	p	t	W	p	t	W	p	t	W	p	
<i>a) Cape knifejaw muscle (n = 5)</i>																						
Percent C	2.23	0.089		-1.31	0.260	15	<b>0.043</b>	-1.02	0.364	4.24	<b>0.013</b>	3.44	<b>0.026</b>	15	<b>0.043</b>	-8.87	<b>0.001</b>	4.29	<b>0.013</b>	-3.45	<b>0.026</b>	
Percent N	-2.08	0.106		-1.64	0.176	3.26	<b>0.031</b>	-13.00	<b>&lt;0.001</b>	2.77	0.051	15	<b>0.043</b>	14	0.080	-14.12	<b>&lt;0.001</b>	1.43	0.227	-4.00	<b>0.016</b>	
C:N atomic	4.81	<b>0.009</b>		2.04	0.111	6.84	<b>0.002</b>	3.96	<b>0.017</b>	4.68	<b>0.009</b>	15	<b>0.043</b>	3.74	<b>0.020</b>	4.49	<b>0.011</b>	7.43	<b>0.002</b>	2.61	0.059	
$\delta^{13}\text{C}$	-1.66	0.172		7.61	<b>0.002</b>	-3.96	<b>0.017</b>	2.03	0.112	1.94	0.125	-2.76	0.051	4.15	0.014	-0.99	0.380	-5.7	<b>0.005</b>	0.29	0.789	
$\delta^{15}\text{N}$	11	0.345		-24.44	<b>&lt;0.001</b>	-4.34	<b>0.012</b>	-4.61	<b>0.010</b>	-6.79	<b>0.002</b>	-49.05	<b>&lt;0.001</b>	-11.22	<b>&lt;0.001</b>	-23.19	<b>&lt;0.001</b>	-0.26	0.806	-19.82	<b>&lt;0.001</b>	
<i>b) Ragged-tooth shark muscle (n = 8)</i>																						
% C	-3.06	<b>0.018</b>		28	<b>0.018</b>	0.11	0.914	-5.64	<b>0.001</b>	30	0.093	36	<b>0.012</b>	-5.33	<b>0.001</b>	36	<b>0.012</b>	36	<b>0.012</b>	-6.31	<b>&lt;0.001</b>	
% N	-1.87	0.104		1.95	0.099	14	1.000	1.38	0.210	4.64	<b>0.002</b>	20	0.779	4.14	<b>0.004</b>	1.65	0.143	33	<b>0.036</b>	2.08	0.076	
C:N atomic	-4.96	<b>0.002</b>		28	<b>0.018</b>	0.94	0.380	-7.79	<b>&lt;0.001</b>	-21.14	<b>&lt;0.001</b>	-6.17	<b>0.001</b>	36	<b>0.012</b>	-12.05	<b>&lt;0.001</b>	36	<b>0.012</b>	-8.95	<b>&lt;0.001</b>	
$\delta^{13}\text{C}$	0.78	0.460		3.53	<b>0.012</b>	1.07	0.321	2.41	<b>0.047</b>	1.73	0.128	19	0.889	2.27	0.058	30	0.093	0.03	0.978	1.84	0.108	
$\delta^{15}\text{N}$	-1.37	0.212		-6.5	<b>0.001</b>	-7.53	<b>&lt;0.001</b>	-5.92	<b>0.001</b>	-5.21	<b>0.001</b>	-14.48	<b>&lt;0.001</b>	-3.43	<b>0.011</b>	-6.01	<b>0.001</b>	-1.54	0.167	-5.77	<b>0.001</b>	
<i>c) Ragged-tooth shark blood (n = 8)</i>																						
% C	-1.31	0.232		36	<b>0.012</b>	6.31	<b>&lt;0.001</b>	36	<b>0.012</b>	5.89	<b>0.001</b>	36	<b>0.012</b>	-13.6	<b>&lt;0.001</b>	36	<b>0.012</b>	-4.26	<b>0.004</b>	36	<b>0.012</b>	
% N	-1.98	0.089		36	<b>0.012</b>	3.49	<b>0.010</b>	8.31	<b>&lt;0.001</b>	36	<b>0.012</b>	6.97	<b>&lt;0.001</b>	36	<b>0.012</b>	10.52	<b>&lt;0.001</b>	36	<b>0.012</b>	36	<b>0.012</b>	
C:N atomic	-0.36	0.727		36	<b>0.012</b>	1.21	0.266	36	<b>0.012</b>	36	<b>0.012</b>	36	<b>0.012</b>	36	<b>0.012</b>	36	<b>0.012</b>	7.13	<b>&lt;0.001</b>	36	<b>0.012</b>	
$\delta^{13}\text{C}$	0.93	0.383		23.69	<b>&lt;0.001</b>	0.34	0.741	16.92	<b>&lt;0.001</b>	14.27	<b>&lt;0.001</b>	12.71	<b>&lt;0.001</b>	22.64	<b>&lt;0.001</b>	13.46	<b>&lt;0.001</b>	6.43	<b>&lt;0.001</b>	18.72	<b>&lt;0.001</b>	
$\delta^{15}\text{N}$	-0.76	0.471		8.93	<b>&lt;0.001</b>	-2.24	0.060	10.16	<b>&lt;0.001</b>	9.41	<b>&lt;0.001</b>	1.65	0.144	10.26	<b>&lt;0.001</b>	7.75	<b>&lt;0.001</b>	-9.83	<b>&lt;0.001</b>	8.89	<b>&lt;0.001</b>	



**FIGURE 3** Effect of subsequent water wash after initial solvent wash on carbon percentage, nitrogen percentage, C:N atomic ratios, and  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values. Significant changes are in black. (Refer to Figure 1 for abbreviations)

Difference among pre-treatments varied with tissues with few pairwise comparisons being significant for  $\delta^{13}\text{C}$  values of ragged-tooth shark muscle, in contrast to the other tissues and  $\delta^{15}\text{N}$  values where most pairwise comparisons between pre-treatments were found to be significant (Table 4, and Tables S1–S3, supporting information). Overall,  $\text{H}_2\text{O}$ -washed muscle samples in both species had significantly higher  $\delta^{15}\text{N}$  values than their counterpart solvent-washed samples (Figure 5). The pre-treatment CM1  $\text{H}_2\text{O}$  exhibited the most different  $\delta^{15}\text{N}$  values in the muscle samples of both species compared with the other 10 pre-treatments. In ragged-tooth shark blood, the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values were most affected by three pre-treatments (i.e. PE, PE  $\text{H}_2\text{O}$ , Cx) which particularly exhibited higher  $\delta^{13}\text{C}$  values than the other eight pre-treatments (Figure 5).

## 4 | DISCUSSION

A review of the literature using SIA to determine the trophic ecology of elasmobranchs highlighted that, despite several studies dealing with pre-treatment issues of lipid and/or urea rich tissues (e.g.<sup>23,25,28</sup>), a consensus has yet to be reached on the preferred pre-treatment. A number of recently published studies used either no pre-treatment (e.g.<sup>8,32–34</sup>), a distilled water rinse followed by a chloroform/methanol 2:1 rinse (e.g.<sup>35</sup>), or cyclohexane rinses (e.g.<sup>36</sup>). The use of different pre-treatments can be problematic with the emergence of studies investigating the trophic ecology of shark species worldwide.<sup>22</sup> Indeed, these pre-treatments may not be equivalent in removing biological compounds such as lipids and urea, and thus they may influence the carbon and nitrogen contents and the C:N ratios,

**TABLE 3** Comparisons across pre-treatments for the five parameters in three tissues. Significant results are in bold


	Cape knifejaw muscle (n = 5)		Ragged-tooth shark muscle (n = 8)		Ragged-tooth shark blood (n = 8)	
	F	p	F	p	F	p
% C	11.47	<b>&lt;0.001</b>	18.28	<b>&lt;0.001</b>	55.48	<b>&lt;0.001</b>
% N	24.35	<b>&lt;0.001</b>	27.82	<b>&lt;0.001</b>	25.15	<b>&lt;0.001</b>
C:N atomic	27.77	<b>&lt;0.001</b>	44.97	<b>&lt;0.001</b>	65.50	<b>&lt;0.001</b>
$\delta^{13}\text{C}$	21.5	<b>&lt;0.001</b>	5.56	<b>&lt;0.001</b>	64.44	<b>&lt;0.001</b>
$\delta^{15}\text{N}$	38.82	<b>&lt;0.001</b>	27.68	<b>&lt;0.001</b>	44.01	<b>&lt;0.001</b>

as well as the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values. While acknowledging the limited sample size, the eleven pre-treatments tested in this study significantly influenced the five measured parameters (carbon and nitrogen percentages, C:N ratios, and  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values; Table 3). The effects were tissue-, species- and pre-treatment-dependent which thus precluded any simple mathematical normalisation.


The differences between pre-treatments are likely to result from a combination of the chemical properties of the rinsing solution and the biological molecules removed (e.g. lipids, urea) that dictate their

solubility. The solutions tested can be placed on a polarity scale with water being the most polar solution followed by chloroform/methanol, chloroform/ethanol, cyclohexane, and petroleum ether in decreasing polarity. Similarly, the polarity of biological compounds depends on their chemical composition and conformation. Urea and TMAO are particularly soluble in polar solutions such as water and less so in solvents such as cyclohexane or petroleum ether.<sup>37,38</sup> On the contrary, lipids in general are insoluble in water but highly soluble in organic solvents.<sup>39</sup> The solubility of lipids in solvents also depends on lipid classes, with neutral lipids such as triacylglycerols being particularly soluble in less polar solvents (e.g. cyclohexane) and phospholipids in more polar solvents (e.g. chloroform).<sup>39</sup>

This suggests that the pre-treatment  $\text{H}_2\text{O}$  removed all water-soluble compounds including urea and TMAO. The removal of urea, depleted in  $^{15}\text{N}$ <sup>40</sup> and probably enriched in  $^{13}\text{C}$ ,<sup>41</sup> would artificially increase  $\delta^{15}\text{N}$  values and potentially lower  $\delta^{13}\text{C}$  values in urea-rich tissues such as in elasmobranchs.<sup>23</sup> The stronger effect of the  $\text{H}_2\text{O}$  rinse was observed in ragged-tooth shark blood for C and N percentages as well as C:N ratios than in the two muscle samples (Figure 2). However, an increase in  $\delta^{15}\text{N}$  values was observed in muscle but not in blood of ragged-tooth sharks where the contrary was observed (Figure 4). Measured urea contents are



**FIGURE 4** Changes in  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values after the eleven pre-treatments relative to the untreated bulk samples of the three tissues (Cape knifejaw muscle, ragged-tooth shark muscle and blood). The data for water-rinsed tissues are in blue. (Refer to Figure 1 for abbreviations) [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 5** Carbon and nitrogen stable isotope ratios (mean  $\pm$  SD) of ragged-tooth shark muscle and blood and Cape knifejaw after water treatment (grey), solvent treatment (black), and solvent and water treatments (dotted line). (Refer to Figure 1 for abbreviations)

relatively homogenous within elasmobranch individuals with very little variation among tissues;<sup>14,42</sup> one would thus expect the effect of urea removal to be similar in blood and muscle. Our results suggest that other water-soluble compounds were removed from the blood concurrently with the urea/TMAO. Elasmobranch blood is characterised by high free amino acid contents compared with muscle.<sup>42</sup> Most of these are soluble in water<sup>37</sup> and would therefore be leached with the  $H_2O$  rinse. Their disappearance would then influence any following chemical analyses. Amino acid contents and compound-specific SIAs would identify whether this amino acid leaching is the reason for differences between blood and muscle in ragged-tooth tissues.

When lipid removal is necessary before SIA, not all lipid extraction protocols are equivalent in terms of quantity and composition of lipids removed (e.g.<sup>43</sup>). Coarsely, lipids can be separated into neutral lipids such as storage lipids (glycerides, wax esters), squalene, sterols, free fatty acids, and structural polar lipids (glycol-, phospho-, sphingo-lipids) which are linked to proteins.<sup>39</sup> Due to the apolarity (or very low polarity) of neutral lipids, they are better extracted with solvents of low polarity (e.g. petroleum ether,

cyclohexane), while structural polar lipids may be extracted with more polar solvents such as chloroform. When removing lipids prior to SIA in lipid-rich tissues, it is preferable to target storage lipids which are highly variable within and among individuals,<sup>44</sup> while preserving the integrity of cell membranes to avoid the co-extraction of associated amino acids. The five solvent mixes tested in this study had varying effects on the five measured parameters (C and N percentages, C:N ratios, and  $\delta^{13}C$  and  $\delta^{15}N$  values). Overall, the least polar solvent (PE) had the least influence on the five parameters measured in the three tissue types (Figures 2 and 4). A PE rinse alone did not change either the  $\delta^{13}C$  or the  $\delta^{15}N$  values in any of the three analysed tissues. Interestingly, the other non-polar solvent, Cx, had the most significant effect on the C percentage and C:N ratios, as well as  $\delta^{13}C$  and  $\delta^{15}N$  values in ragged-tooth shark blood. A direct influence of Cx carbons into the results can be ruled out as this pattern is not observed in muscle tissues. These results suggest that there may be a carbon-rich molecule/component present in blood but not in muscle, and insoluble in PE, that could affect the five measured parameters following a Cx extraction. The identity of this molecule is currently unknown and would require further research.

Instead of using chemicals, mathematical normalisations are sometimes used after a  $H_2O$  rinse to remove the impact of lipids on  $\delta^{13}C$  values.<sup>45</sup> Several mathematical models exist (e.g.<sup>19,46,47</sup>) and they are not all equivalent.<sup>48</sup> When testing two widely used models developed using multispecies data<sup>19,46</sup> with our results, the models consistently predicted higher  $\delta^{13}C$  values (up to 2‰) for all three tissues and solvents tested except for whole blood rinsed with PE and Cx where the  $\delta^{13}C$  values were lower than expected with both models (-0.6‰). As pointed out previously (e.g.<sup>46,48,49</sup>), this stresses the need for species- and tissue-specific equations developed on a sub-set of samples. Importantly, our work also shows that these equations will also depend on the chemicals used to remove the lipids as not all pre-treatments are equivalent (see above).

The comparison of pre-treatments consisting of the solvent rinse followed by the water rinse showed that this approach reduced the difference among treatments, particularly with regard to the C and N percentages in blood. The comparison of  $\delta^{13}C$  and  $\delta^{15}N$  values among pre-treatments showed that the measured differences were higher than any instrument errors for all three tissues (Figures 4 and 5). The differences in  $\delta^{13}C$  values between the various pre-treatments may be as high as 3‰, but most often are about 1‰ (Figure 4). In both ragged-tooth shark tissues, the corrected  $\delta^{13}C$  values were  $\sim$ 1‰ higher than those in bulk tissues, while the contrary was true for Cape knifejaw muscle.  $\delta^{13}C$  values provide insight into the source production at the base of the food chain,<sup>50</sup> and modelling studies using isoscapes to retrospectively locate predators are multiplying (e.g.<sup>51</sup>). An artificial variation of  $\delta^{13}C$  value due to laboratory protocols may therefore lead to erroneous ecological interpretation. The variability among pre-treatments was higher for  $\delta^{15}N$  values. The most notable impact was observed in  $\delta^{15}N$  values of ragged-tooth shark muscle with differences of up to 3‰ (i.e. a trophic level<sup>52</sup>) between PE and CM1 treatments. With regard to the results from the other two tissues, the effects of the pre-treatments resulted in significant differences of about 2‰.

The range of these differences is likely to influence ecological interpretations in global studies, particularly when  $\delta^{15}\text{N}$  values rather than trophic levels are compared.

In addition to the influence of pre-treatments on the five parameters measured, the variability among samples within the three sample types was also noticeable (Figures 2 and 4). Patterns of enrichment were particularly variable for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values (Figure 4). This was probably linked to variable urea, TMAO and lipid contents in the individual fish analysed. For example, as all the sharks were captured over a period of 18 months, they were variable in length (188 to 287 cm total length; Smale, unpublished data) and sex (2 males, 4 females; Smale, unpublished data), and these factors may have contributed to differing quantities of urea, TMAO and lipid being present in the tissues. This would have influenced how much of these compounds was removed by the pre-treatments and thus introduced variability in the impact of these pre-treatments on the corrected values.

Finally, the high C:N ratios observed in Cape knifejaw muscle ( $\text{C:N} \geq 4.1$ ) indicated that lipid extraction had to be undertaken to remove the lowering effect of lipids on  $\delta^{13}\text{C}$  value.<sup>19,20</sup> Initial low C:N ratios observed in ragged-tooth shark tissues ( $\text{C:N} < 3.5$ ) suggested that a lipid extraction was not necessary. However, after removing water-soluble compounds, the C:N ratios increased to above 4 in both shark tissues, suggesting a potential confounding effect of lipids on the  $\delta^{13}\text{C}$  values. Such results indicate that care must be taken to ensure that the most appropriate pre-treatment is used prior to SIA. The lipid content of muscle tissue from ragged-tooth sharks is particularly high in comparison with that of other shark species found in South African waters.<sup>53</sup> Our study confirms the inadequacy of using raw C:N ratios as an indication of lipid contents in elasmobranchs.<sup>21,25</sup>

## 5 | CONCLUDING REMARKS

Although lipid and urea removals are used in most food web studies including elasmobranchs, the various pre-treatments yield statistically significant different results.

- Caution is therefore needed when using SI values from the literature because the pre-treatments vary widely between studies; the compared SI data thus reflect the intrinsic differences in the tissue composition combined with the influence of the pre-treatments which may confound ecological interpretation;
- Our study, conducted on a limited number of samples, shows that the difference among pre-treatments extends beyond the elasmobranch species and should therefore be confirmed with tests on additional species;
- The pre-treatment comparative approach that we followed for ragged-tooth sharks and Cape knifejaws could be adapted for other species prior to large scale studies to provide conversion factors;
- The next step would entail the evaluation of the pre-treatments in their abilities in removing solely the biological molecules of interest (urea, TMAO, lipids). This would allow us to choose the

best pre-treatment which interferes least with the amino acid composition;

- We reiterate the call for a standardised approach to pre-treatments which would greatly benefit larger-scale comparisons of SI studies, particularly when global comparisons of wide-ranging species are starting to emerge (e.g.<sup>22</sup>).

## ACKNOWLEDGEMENTS

The authors would like to thank Dr D. Zimmerman (National Parks Board) for his veterinary expertise, field participation and oversight of the study, and M. Bradshaw for her assistance during fieldwork, sample preparation and storage. Mr M. McLean is thanked for skipping the boat during research tagging operations. Field work was funded by the African Coelacanth Ecosystem Programme of the National Research Foundation (Grant No. 81879).

## ORCID

Maëlle Connan  <https://orcid.org/0000-0002-1308-9118>

Grant Hall  <https://orcid.org/0000-0002-2164-4900>

Malcolm Smale  <https://orcid.org/0000-0002-1135-8687>

## REFERENCES

1. Hobson KA. Tracing origins and migration of wildlife using stable isotopes: a review. *Oecologia*. 1999;120(3):314-326. <https://doi.org/10.1007/s004420050865>
2. Chelou Y, Hobson KA, Bailleul F, Groscolas R. Nutrition, physiology, and stable isotopes: new information from fasting and molting penguins. *Ecology*. 2005;86(11):2881-2888. <https://doi.org/10.1890/05-0562>
3. Roff G, Doropoulos C, Rogers A, et al. The ecological role of sharks on coral reefs. *Trends Ecol Evol*. 2016;31(5):395-407. <https://doi.org/10.1016/j.tree.2016.02.014>
4. Hoffman JC. Tracing the origins, migrations, and other movements of fishes using stable isotopes. In: Morais P, Daverat F, eds. *An Introduction to Fish Migration*. Boca Raton: CRC Press; 2016:169-196.
5. Gannes LZ, Martínez del Río C, Koch P. Natural abundance variations in stable isotopes and their potential uses in animal physiological ecology. *Comp Biochem Physiol - Part A*. 1998;119(3):725-737. [https://doi.org/10.1016/S1095-6433\(98\)01016-2](https://doi.org/10.1016/S1095-6433(98)01016-2)
6. Hussey NE, Dudley SFJ, McCarthy ID, Cliff G, Fisk AT. Stable isotope profiles of large marine predators: viable indicators of trophic position, diet and movement in sharks. *Can J Fish Aquat Sci*. 2011;68(12):2029-2045. <https://doi.org/10.1139/f2011-115>
7. Dicken ML, Hussey NE, Christiansen HM, et al. Diet and trophic ecology of the tiger shark (*Galeocerdo cuvier*) from South African waters. *PLoS One*. 2017;12(6):e0177897. <https://doi.org/10.1371/journal.pone.0177897>
8. Ferreira LC, Thums M, Heithaus MR, et al. The trophic role of a large marine predator, the tiger shark *Galeocerdo cuvier*. *Sci Rep*. 2017;7(1):7641. <https://doi.org/10.1038/s41598-017-07751-2>
9. Rosas-Luis R, Navarro J, Lóor-Andrade P, Forero MG. Feeding ecology and trophic relationships of pelagic sharks and billfishes coexisting in the central eastern Pacific Ocean. *Mar Ecol Prog Ser*. 2017;573:191-201. <https://doi.org/10.3354/meps12186>
10. Dulvy NK, Baum JK, Clarke S, et al. You can swim but you can't hide: the global status and conservation of oceanic pelagic sharks and rays. *Aquat Conserv*. 2008;18(5):459-482. <https://doi.org/10.1002/aqc.975>
11. Dalerum F, Angerbjörn A. Resolving temporal variation in vertebrate diets using naturally occurring stable isotopes. *Oecologia*. 2005;144(4):647-658. <https://doi.org/10.1007/s00442-005-0118-0>

12. Buchheister A, Latour RJ. Turnover and fractionation of carbon and nitrogen stable isotopes in tissues of a migratory coastal predator, summer flounder (*Paralichthys dentatus*). *Can J Fish Aquat Sci.* 2010;67(3):445-461. <https://doi.org/10.1139/F09-196>
13. Goldstein L. Urea biosynthesis in elasmobranchs. In: Gilbert PW, Mathewson RF, Rall DP, eds. *Sharks, Skates and Rays*. Baltimore: Johns Hopkins University Press; 1967:207-214.
14. Ballantyne JS. Jaws: The inside story. The metabolism of elasmobranch fishes. *Comp Biochem Physiol - Part B.* 1997;118(4):703-742. [https://doi.org/10.1016/S0305-0491\(97\)00272-1](https://doi.org/10.1016/S0305-0491(97)00272-1)
15. Hazon N, Wells A, Pillard RD, Good JP, Anderson WG, Franklin CE. Urea based osmoregulation and endocrine control in elasmobranch fish with special reference to euryhalinity. *Comp Biochem Physiol - Part B.* 2003;136(4):685-700. [https://doi.org/10.1016/S1096-4959\(03\)00280-X](https://doi.org/10.1016/S1096-4959(03)00280-X)
16. Steele KW, Daniel RM. Fractionation of nitrogen isotopes by animals: a further complication to the use of variations in the natural abundance of  $^{15}\text{N}$  for tracer studies. *J Agric Sci.* 1978;90(01):7-9.
17. DeNiro MJ, Epstein S. Mechanism of carbon isotope fractionation associated with lipid synthesis. *Science.* 1977;197(4300):261-263. <https://doi.org/10.1126/science.327543>
18. McConnaughey T, McRoy CP. Food-web structure and the fractionation of carbon isotopes in the Bering Sea. *Mar Biol.* 1979;53(3):257-262. <https://doi.org/10.1007/BF00952434>
19. Post DM, Layman CA, Arrington DA, Takimoto G, Quattrochi J, Montaña CG. Getting to the fat of the matter: models, methods and assumptions for dealing with lipids in stable isotope analyses. *Oecologia.* 2007;152(1):179-189. <https://doi.org/10.1007/s00442-006-0630-x>
20. Skinner MM, Martin AA, Moore BC. Is lipid correction necessary in the stable isotope analysis of fish tissues? *Rapid Commun Mass Spectrom.* 2016;30(7):881-889. <https://doi.org/10.1002/rcm.7480>
21. Carlisle AB, Litvin SY, Madigan DJ, et al. Interactive effects of urea and lipid content confound stable isotope analysis in elasmobranch fishes. *Can J Fish Aquat Sci.* 2017;74(3):419-428. <https://doi.org/10.1139/cjfas-2015-0584>
22. Bird CS, Verissimo A, Magozzi S, et al. A global perspective on the trophic geography of sharks. *Nature Ecol Evol.* 2018;2(2):299-305. <https://doi.org/10.1038/s41559-017-0432-z>
23. Kim SL, Koch PL. Methods to collect, preserve, and prepare elasmobranch tissues for stable isotope analysis. *Environ Biol Fishes.* 2012;95(1):53-63. <https://doi.org/10.1007/s10641-011-9860-9>
24. Kiszka JJ, Charlot K, Hussey NE, et al. Trophic ecology of common elasmobranchs exploited by artisanal shark fisheries off south-western Madagascar. *Aquat Biol.* 2014;23(1):29-38. <https://doi.org/10.3354/ab00602>
25. Li Y, Zhang YP, Hussey NE, Dai X. Urea and lipid extraction treatment effects on  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values in pelagic sharks. *Rapid Commun Mass Spectrom.* 2016;30(1):1-8. <https://doi.org/10.1002/rcm.7396>
26. Hussey NE, Brush J, McCarthy ID, Fisk AT.  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  diet-tissue discrimination factors for large sharks under semi-controlled conditions. *Comp Biochem Physiol - Part A.* 2010;155(4):445-453. <https://doi.org/10.1016/j.cbpa.2009.09.023>
27. Hussey NE, MacNeil MA, Olin JA, et al. Stable isotopes and elasmobranchs: tissue types, methods, applications and assumptions. *J Fish Biol.* 2012;80(5):1449-1484. <https://doi.org/10.1111/j.1095-8649.2012.03251.x>
28. Hussey NE, Olin JA, Kinney MJ, McMeans BC, Fisk AT. Lipid extraction effects on stable isotope values ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) of elasmobranch muscle tissue. *J Exp Mar Biol Ecol.* 2012;434-435:7-15. <https://doi.org/10.1016/j.jembe.2012.07.012>
29. Woodborne S, Huchzermeyer KDA, Govender D, et al. Ecosystem change and the Olifants River crocodile mass mortality events. *Ecosphere.* 2012;3(10):1-17. <https://doi.org/10.1890/ES12-00170.1>
30. Hammer Ø, Harper DAT, Ryan PD. PAST: Palaeontological Statistics software package for education and data analysis. *Palaeontologia Electronica.* 2001;4(1):9p.
31. R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing; 2018.
32. Gallagher AJ, Shiffman DS, Byrnes EE, Hammerschlag-Peyer CM, Hammerschlag N. Patterns of resource use and isotopic niche overlap among three species of sharks occurring within a protected subtropical estuary. *Aquat Ecol.* 2017;51(3):435-448. <https://doi.org/10.1007/s10452-017-9627-2>
33. Matich P, Kiszka JJ, Mourier J, Planes S, Heithaus M. Species co-occurrence affects the trophic interactions of two juvenile reef shark species in tropical lagoon nurseries in Moorea (French Polynesia). *Mar Environ Res.* 2017;127:84-91. <https://doi.org/10.1016/j.marenvres.2017.03.010>
34. Barria C, Navarro J, Coll M. Trophic habits of an abundant shark in the northwestern Mediterranean Sea using an isotopic non-lethal approach. *Estuar Coast Shelf Sci.* 2018;207:383-390. <https://doi.org/10.1016/j.ecss.2017.08.021>
35. Poulakis GR, Urakawa H, Stevens PW, et al. Sympatric elasmobranchs and fecal samples provide insight into the trophic ecology of the smalltooth sawfish. *Endanger Species Res.* 2017;32:491-506. <https://doi.org/10.3354/esr00824>
36. Kiszka JJ, Aubail A, Hussey NE, Heithaus MR, Caurant F, Bustamante P. Plasticity of trophic interactions among sharks from the oceanic south-western Indian Ocean revealed by stable isotope and mercury analyses. *Deep-Sea Research I.* 2015;96:49-58. <https://doi.org/10.1016/j.dsr.2014.11.006>
37. Budavari S. *The Merck Index - An Encyclopedia of Chemicals, Drugs, and Biologicals*. Whitehouse Station, NJ: Merck and Co., Inc.; 1996.
38. Wishart DS, Feunang YD, Marcu A, et al. HMDB 4.0 - The Human Metabolome Database for 2018. *Nucleic Acids Res.* 2018;46(D1):D608-D617. Available at <http://www.hmdb.ca/> accessed 6 September 2018
39. Christie WW. *Advances in Lipid Methodology - Two*. Dundee: Oily Press; 1993.
40. Fisk AT, Tittlemier SA, Pranschke JL, Norstrom RJ. Using anthropogenic contaminants and stable isotopes to assess the feeding ecology of Greenland sharks. *Ecology.* 2002;83(8):2162-2172. <https://doi.org/10.2307/3072048>
41. Ilev A, Knyazev YA, Logachev M. Daily average carbon isotope composition of  $\text{CO}_2$  of expired air and urine urea in norm and some endocrine pathologies in man. *Biofizika.* 1996;41(2):508-516.
42. Bedford JJ. The composition of the fluid compartments of two chondrichthyans, *Callorhynchus milli* and *Squalus acanthias*. *Comp Biochem Physiol - Part A.* 1983;76(1):75-80. [https://doi.org/10.1016/0300-9629\(83\)90295-5](https://doi.org/10.1016/0300-9629(83)90295-5)
43. Iverson SJ, Lang SLC, Cooper MH. Comparison of the Bligh and Dyer and Folch methods for total lipid determination in a broad range of marine tissue. *Lipids.* 2001;36(11):1283-1287.
44. Pethybridge H, Daley R, Virtue P, Nichols P. Lipid composition and partitioning of deepwater chondrichthyans: inferences of feeding ecology and distribution. *Mar Biol.* 2010;157(6):1367-1384. <https://doi.org/10.1007/s00227-010-1416-6>
45. Torres Rojas YE, Osuna FP, Herrera AH, et al. Feeding grounds of juvenile scalloped hammerhead sharks (*Sphyrna lewini*) in the south-eastern Gulf of California. *Hydrobiologia.* 2014;726(1):81-94. <https://doi.org/10.1007/s10750-013-1753-9>
46. Kiljunen M, Grey J, Sinisalo T, Harrod C, Immonen H, Jones RI. A revised model for lipid-normalizing  $\delta^{13}\text{C}$  values from aquatic organisms, with implications for isotope mixing models. *J Appl Ecol.* 2006;43(6):1213-1222. <https://doi.org/10.1111/j.1365-2664.2006.01224.x>
47. Hoffman JC, Sutton TT. Lipid correction for carbon stable isotope analysis of deep-sea fishes. *Deep-Sea Research I.* 2010;57(8):956-964. <https://doi.org/10.1016/j.dsr.2010.05.003>

48. Reum JCP. Lipid correction model of carbon stable isotopes for a cosmopolitan predator, spiny dogfish *Squalus acanthias*. *J Fish Biol.* 2011;79(7):2060-2066. <https://doi.org/10.1111/j.1095-8649.2011.03120.x>
49. Logan JM, Jardine TD, Miller TJ, Bunn SE, Cunjak RA, Lutcavage ME. Lipid corrections in carbon and nitrogen stable isotope analyses: comparison of chemical extraction and modelling methods. *J Anim Ecol.* 2008;77(4):838-846.
50. Michener RH, Kaufman L. Stable isotope ratios as tracers in marine food webs: An update. In: Michener R, Lajtha K, eds. *Stable isotopes in ecology and environmental science*. Singapore: Blackwell Publishing Ltd; 2007:238-282.
51. Trueman CN, MacKenzie KM, Palmer MR. Identifying migrations in marine fishes through stable-isotope analysis. *J Fish Biol.* 2012;81(2):826-847. <https://doi.org/10.1111/j.1095-8649.2012.03361.x>
52. Caut S, Angulo E, Courchamp F. Variation in discrimination factors ( $\Delta^{15}\text{N}$  and  $\Delta^{13}\text{C}$ ): the effect of diet isotopic values and applications for diet reconstruction. *J Appl Ecol.* 2009;46(2):443-453. <https://doi.org/10.1111/j.1365-2664.2009.01620.x>
53. Davidson B, Sidell J, Rhodes J, Cliff G. A comparison of the heart and muscle total lipid and fatty acid profiles of nine large shark species from the east coast of South Africa. *Fish Physiol Biochem.* 2011;37(1):105-112. <https://doi.org/10.1007/s10695-010-9421-8>

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Connan M, Hall G, Smale M. Effects of pre-treatments on bulk stable isotope ratios in fish samples: A cautionary note for studies comparisons. *Rapid Commun Mass Spectrom.* 2019;33:291-302. <https://doi.org/10.1002/rcm.8344>